

Each of the following are depictions of the **same** relation, the **dept** relation, whose relation structure form can be given as:
dept(dept_name, DEPT_NUM, dept_loc)

dept_name dept_num dept_loc

Accounting	10	New York
Research	20	Dallas
Sales	30	Chicago
Operations	40	Boston

dept_num dept_name dept_loc

40	Operations	Boston
30	Sales	Chicago
20	Research	Dallas
10	Accounting	New York

Relational selection example

For example, say that you have a **Student** relation as follows:

STUDENT table:

Stud_ ID	Stud_Name	Stud_Major	Stud_Grade _Level	Stud_Age
123	Jones	History	JR	21
158	Parks	Math	GR	26
105	Anderson	Management	SR	27
271	Smith	History	JR	19

Then the result of the **selection** operation on the **Student** table of rows in which **Age < 25** would be:

STUDENT WHERE Age < 25

Stud_ ID	Stud_Name	Stud_Major	Stud_Grade _Level	Stud_Age
123	Jones	History	JR	21
271	Smith	History	JR	19

Relational projection example

For example, say that you have a **Student** relation as follows:

STUDENT table:

Stud_ ID	Stud_Name	Stud_Major	Stud_Grade _Level	Stud_Age
123	Jones	History	JR	21
158	Parks	Math	GR	26
105	Anderson	Management	SR	27
271	Smith	History	JR	19

Then the result of the **projection** operation of the **Student_Major** and **Stud_grade_level** attributes of the **Student** table would be:

Stud_Major	Stud_Grade_Level
History	JR
Math	GR
Management	SR

Relational Cartesian product example

For example, say that you have a **Price** relation as follows:

Prod_Code	Price
AA	5.99
BB	22.75

And, say that you have a **Location** table as follows:

Store	Aisle	Shelf
23	W	5
24	K	9
25	Z	6

Then the **Cartesian product** of Price and Location would be:

Prod_Code	Price	Store	Aisle	Shelf
AA	5.99	23	W	5
AA	5.99	24	K	9
AA	5.99	25	Z	6
BB	22.75	23	W	5
BB	22.75	24	K	9
BB	22.75	25	Z	6

Relational equi-join and natural join examples

For example, say that you have **Student** and **Enrollment** tables as follows:

STUDENT table:

Stud_ ID	Stud_Name	Stud_Major	Stud_Grade_ Level	Stud_Age
123	Jones	History	JR	21
158	Parks	Math	GR	26
105	Anderson	Management	SR	27
271	Smith	History	JR	19

ENROLLMENT table:

Stud_ID	Class_Name	Position_Num
123	H350	1
105	BA490	3
123	BA490	7

Say that you wish to compute the **equi-join** and **natural join** of these tables based on the join condition (**Student.stud_id = Enrollment.stud_id**).

First, compute the **Cartesian product** of these two tables:

Student. Stud_ID	Stud_ Name	Stud_ Major	Stud_ Grade_ Level	Stud_ Age	Enrollment .Stud_ID	Class_ Name	Position _Num
123	Jones	History	JR	21	123	H350	1
123	Jones	History	JR	21	105	BA490	3
123	Jones	History	JR	21	123	BA490	7
158	Parks	Math	GR	26	123	H350	1
158	Parks	Math	GR	26	105	BA490	3
158	Parks	Math	GR	26	123	BA490	7
105	Anderson	Management	SR	27	123	H350	1
105	Anderson	Management	SR	27	105	BA490	3
105	Anderson	Management	SR	27	123	BA490	7
271	Smith	History	JR	19	123	H350	1
271	Smith	History	JR	19	105	BA490	3
271	Smith	History	JR	19	123	BA490	7

Second, perform a selection on this result of only those rows for which (**Student.stud_id = Enrollment.stud_id**):

Student. Stud_ID	Stud_ Name	Stud_ Major	Stud_ Grade_ Level	Stud_ Age	Enrollment .Stud_ID	Class_ Name	Position _Num
123	Jones	History	JR	21	123	H350	1
123	Jones	History	JR	21	123	BA490	7
105	Anderson	Management	SR	27	105	BA490	3

This is the **equi-join** of these two tables on the join condition (**Student.stud_id = Enrollment.stud_id**).

Then, the **natural join** of these two tables on this join condition would include the **third** step of now projecting all of the columns in this result except for one of the "duplicate-contents" columns (and it doesn't matter which one of the two is omitted).

So, for example, the resulting **natural join** in this case could be:

Stud_ID	Stud_ Name	Stud_ Major	Stud_ Grade_ Level	Stud_ Age	Class_ Name	Position_ Num
123	Jones	History	JR	21	H350	1
123	Jones	History	JR	21	BA490	7
105	Anderson	Management	SR	27	BA490	3